

A Unified View on the Functorial Nerve Theorem and its Variations

Fabian Roll (TUM)



Group meeting - NDAG
September 8, 2022

joint work with Ulrich Bauer, Michael Kerber, and Alexander Rolle

The Alexandrov nerve (1928)

Definition. Let X be a topological space, and let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of X . The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid |J| < \infty \text{ and } U_J := \bigcap_{i \in J} U_i \neq \emptyset\}$$




The Alexandrov nerve (1928)

Čech complex of a point cloud

Definition. The Čech complex of a subset $X \subseteq \mathbb{R}^d$ is the nerve of the cover by closed balls of radius r centered at points in X

$$\check{C}ech_r(X) = \text{Nrv}((D_r(X))_{x \in X})$$



The history of the nerve theorem

Theorem (Borsuk 1948, and many more). Let \mathcal{U} be a nice cover of X . Then $\text{Nrv}(\mathcal{U})$ is homotopy equivalent to X .

Here *nice* can mean different things:

- ▶ open, numerable, and good cover (contractible intersections)
→ many references
- ▶ closed, finite, and convex cover
→ few references, mostly using outdated language and tools

Prior results?


- ▶ Alexandrov 1928: Every compact metric space is the inverse limit of a sequence of nerves of “arbitrarily fine” closed covers.
- ▶ Čech 1932: Extends Alexandrov’s ideas → Čech (co)homology

Nerve theorem for closed convex covers

Theorem. If $X \subset \mathbb{R}^d$, and $\mathcal{A} = (C_i)_{i \in [n]}$ is a cover by closed convex subsets, then $\text{Nrv}(\mathcal{A})$ is homotopy equivalent to X .

Our proof strategy:

- ▶ Construct piecewise linear $\Gamma: \text{Sd Nrv}(\mathcal{A}) \rightarrow X$ with $\Gamma(\text{bst } v_i) \subseteq C_i$.



- ▶ Construct $\Phi: X \rightarrow \text{Nrv}(\mathcal{A})$ using a partition of unity subordinate to an open thickening of the C_i with $\Phi(C_i) \subseteq \text{bst } v_i$.
- ▶ Show that Φ is a homotopy inverse to Γ .

Nerve theorem for closed convex covers

Some proof details

- ▶ Take a collection of open sets $\mathcal{G} = (U_i \supseteq C_i)_{i \in [n]}$ satisfying $\text{Nrv}(\mathcal{A}) = \text{Nrv}(\mathcal{G}_\varepsilon)$.
- ▶ Take a partition of unity $(\psi_i)_{i \in [n]}$ on X subordinate to the cover $(U_i \cap X)_{i \in [n]}$ of X with $\psi_i|_{C_i} \equiv 1$.
- ▶ Define the map $\Phi: X \rightarrow \text{Nrv}(\mathcal{A})$ by

$$\Phi: x \mapsto \sum_{i=0}^n \psi_i(x) \cdot v_i$$

- ▶ Then $\Gamma \circ \Phi(C_i) \subseteq C_i$ and $\Gamma \circ \Phi \simeq \text{id}_X$ by a straight line homotopy.
- ▶ Then $\Phi \circ \Gamma(\text{bst } v_i) \subseteq \text{bst } v_i$ and $\Phi \circ \Gamma \simeq \text{id}_{\text{Sd Nrv}(\mathcal{A})}$ by induction over the skeleton of $\text{Sd Nrv}(\mathcal{A})$; use that $(\text{bst } v)_{v \in \text{Vert Nrv}(\mathcal{A})}$ is a good cover of $\text{Sd Nrv}(\mathcal{A})$.

Nerve theorem for simplicial covers

Theorem (Björner, Quillen 1979). If $\mathcal{A} = (K_i \subseteq K)_{i \in I}$ is a good cover by subcomplexes, then $\text{Nrv}(\mathcal{A})$ is homotopy equivalent to K .

Proof strategy:

- ▶ Use Quillen's Theorem A: If $f: P \rightarrow Q$ is a map of posets, and y/f is contractible for all $y \in Q$, then f is a homotopy equivalence.
- ▶ Consider the product poset

$$Z = \{(\sigma, J) \mid \sigma \in K_J\} \subseteq \text{Pos}(K) \times \text{Pos}(\text{Nrv}(\mathcal{A}))$$


and apply Quillen's Theorem A to the projections

$$\text{Pos}(K) \longleftarrow Z \longrightarrow \text{Pos}(\text{Nrv}(\mathcal{A})) \quad \text{op}$$


locally finite cover

Functoriality

Persistent homology



Functoriality

Let $X \subseteq \mathbb{R}^d$, $\mathcal{U}_r = \{D_r(x)\}_{x \in X}$, and $X_r = \bigcup \mathcal{U}_r$. The nerve theorem guarantees for $r \leq l$ the homotopy equivalences

$$\begin{array}{ccc} \mathrm{Nrv}(\mathcal{U}_r) & \longrightarrow & \mathrm{Nrv}(\mathcal{U}_l) \\ \simeq \uparrow & \circlearrowleft? & \uparrow \simeq \\ X_r & \longrightarrow & X_l \end{array}$$


Theorem (Chazal–Oudot 2008). For **open covers**, this diagram commutes **up to homotopy**. Hence, it commutes after applying homology.

We address two issues:

1. Closed covers were not well-treated in the literature
2. No “proper” functoriality \rightarrow needed in some homotopy-theoretic approaches to TDA (e.g. Blumberg–Lesnick)

Functoriality

Category of covered spaces



Definition. $(U_i)_{i \in I}$ a cover of X , and $(V_\ell)_{\ell \in L}$ a cover of Y . A *map of indexed covers* $\varphi: (U_i)_{i \in I} \rightarrow (V_\ell)_{\ell \in L}$ is formally a map $\varphi: I \rightarrow L$. A continuous map $f: X \rightarrow Y$ is *carried by* φ if $f(U_i) \subseteq V_{\varphi(i)}$.

Definition. The *category of covered spaces* Cov has

- ▶ Obj: pairs of the form $(X, (U_i))$, with (U_i) a cover of X
- ▶ Mor: $(f, \varphi): (X, (U_i)) \rightarrow (Y, (V_\ell))$, continuous map $f: X \rightarrow Y$ carried by $\varphi: (U_i) \rightarrow (V_\ell)$


Functoriality

Category of covered spaces

Two functors

- ▶ Forgetting the cover: $\text{Spc}: \text{Cov} \rightarrow \text{Top}, (X, \mathcal{U}) \mapsto X$
- ▶ The nerve: $\text{Nrv}: \text{Cov} \rightarrow \text{Top}, (X, \mathcal{U}) \mapsto \text{Nrv}(\mathcal{U})$


Remark. There are no natural transformations $\text{Spc} \Rightarrow \text{Nrv}$



and similarly no natural transformations $\text{Nrv} \Rightarrow \text{Spc}$.

Functoriality

Pointed covers



Definition. Only consider $X \subseteq \mathbb{R}^d$. The category ClConv_\bullet has

- ▶ Obj: (X, \mathcal{A}_\bullet) , \mathcal{A}_\bullet a finite closed, convex, and *pointed* cover of X
- ▶ Mor: $(f, \varphi): (X, \mathcal{A}_\bullet) \rightarrow (Y, \mathcal{B}_\bullet)$, $f: X \rightarrow Y$ carried by $\varphi: \mathcal{A} \rightarrow \mathcal{B}$:
 - ▶ f preserves the basepoints
 - ▶ f is affine linear on each cover element

The map $\Gamma: \text{Sd Nrv}(\mathcal{A}) \xrightarrow{\cong} X$ is natural w.r.t morphisms in ClConv_\bullet .

Theorem. On ClConv_\bullet there exists a pointwise homotopy equivalence

$$\text{Sd Nrv} \Rightarrow \text{Spc}$$



Functoriality

Blowup complex

Definition. For a finite cover $\mathcal{U} = (U_i)_{i \in [n]}$ of X , the *blowup complex* is

$$\text{Blowup}(\mathcal{U}) = \bigcup_{J \in \text{Nrv}(\mathcal{U})} U_J \times \Delta^{|J|-1} \subseteq X \times \Delta^n,$$


yielding a functor $\text{Blowup}: \text{Cov} \rightarrow \text{Top}$.



Functoriality

Bar construction

Remark. The blowup complex is (not naturally) homeomorphic to the *bar construction* of the nerve diagram



→ can exploit its good homotopical properties to prove nerve theorems

Functoriality

Any $(f, \varphi): (X, \mathcal{U}) \rightarrow (Y, \mathcal{V})$ induces a commuting diagram

$$\begin{array}{ccccc} X & \xleftarrow{\rho_S} & \text{Blowup}(\mathcal{U}) & \xrightarrow{\rho_N} & \text{Nrv}(\mathcal{U}) \\ f \downarrow & & \downarrow & & \downarrow \varphi_* \\ Y & \xleftarrow{\rho_S} & \text{Blowup}(\mathcal{V}) & \xrightarrow{\rho_N} & \text{Nrv}(\mathcal{V}) \end{array}$$

Hence, there are natural transformations $\text{Spc} \xleftarrow{\rho_S} \text{Blowup} \xrightarrow{\rho_N} \text{Nrv}$.

Theorem. If $X \subset \mathbb{R}^d$, and $\mathcal{A} = (C_i)_{i \in [n]}$ is a cover by closed convex subsets, then the natural maps ρ_S and ρ_N are homotopy equivalences.

Proof. Use the following up to homotopy commutative diagram

$$\begin{array}{ccc} & \text{Blowup}(\mathcal{A}) & \\ \nearrow \simeq & & \searrow \simeq \\ X & \xrightarrow[\Gamma]{\simeq} & \text{Nrv}(\mathcal{A}) \end{array}$$

Functoriality

Simplicial covers

Definition. For a cover $\mathcal{A} = (K_i \subseteq K)_{i \in I}$ by subcomplexes, consider the subset

$$\text{PoBar}(\mathcal{A}) = \{(\sigma, J) \mid J \subseteq I \text{ finite}, \sigma \in K_J\}$$

of $\text{Pos}(K) \times \text{Pos}(\text{Nrv}(\mathcal{A}))^{\text{op}} \rightarrow \text{Bar}$ construction in the cat. of posets!

Remark. There exists a commuting diagram

$$\begin{array}{ccccc} K & \xleftarrow{\rho_S} & \text{Blowup}(\mathcal{A}) & \xrightarrow{\rho_N} & \text{Nrv}(\mathcal{A}) \\ \cong \downarrow & & \downarrow \cong & & \downarrow \cong \\ \text{Sd}(K) & \xleftarrow{\quad} & \text{T}(\mathcal{A}) & \xrightarrow{\quad} & \text{Sd}(\text{Nrv}(\mathcal{A})) \\ & \swarrow \lambda_S & \downarrow \text{red} & \searrow \lambda_N & \\ & & \text{Flag}(\text{PoBar}(\mathcal{A})) & & \end{array}$$

Unified nerve theorem

Theorem. Let X be a topological space and $\mathcal{A} = (A_i)_{i \in I}$ a cover of X .

1. Consider the natural map $\rho_S: \text{Blowup}(\mathcal{A}) \rightarrow X$.
 - a) If \mathcal{A} is an open cover, then ρ_S is a weak homotopy equivalence. If furthermore X is paracompact, then ρ_S is a homotopy equivalence.
 - b) Let X be compactly generated, and \mathcal{A} a closed cover that is locally finite and locally finite dimensional. If for any $T \in \text{Nrv}(\mathcal{A})$ the latching space $L(T) := \bigcup_{T \subsetneq J \subseteq I} A_J \subseteq A_T$ is closed and $(A_T, L(T))$ satisfies the homotopy ext. prop., then ρ_S is a homotopy equivalence.
2. Consider the natural map $\rho_N: \text{Blowup}(\mathcal{A}) \rightarrow \text{Nrv}(\mathcal{A})$.
 - a) If \mathcal{A} is (weakly) good, then ρ_N is a (weak) homotopy equivalence.
 - b) If for all $J \in \text{Nrv}(\mathcal{A})$ the space A_J is compactly generated and \mathcal{A} is homologically good with respect to a coefficient ring R , then ρ_N is an R -homology isomorphism.

Unified nerve theorem

Counterexamples

1. Paracompactness is crucial to guarantee a homotopy equivalence
2. For closed good covers some “finiteness” is needed
3. The “latching assumption” is not a proof artefact; even if we only care about the homologies

Unified nerve theorem

Counterexamples

1. Paracompactness is crucial to guarantee a homotopy equivalence:
 - ▶ Consider the *long ray* L
 - ▶ This is a standard example for a non-paracompact space that is also not contractible
 - ▶ L is weakly contractible and for any point $p \in L$ the open set $L_{<p} = \{t \in L \mid t < p\}$ is homeomorphic to the interval $[0, 1)$
 - ▶ $\mathcal{A} = (L_{<p})_{p \in \omega_1}$ is a good open cover and it follows from 1.a) and 2.a) that the nerve $\text{Nrv } \mathcal{A}$ is weakly contractible and hence contractible by Whitehead's theorem
 - ▶ Thus, L and $\text{Nrv } \mathcal{A}$ are not homotopy equivalent

Unified nerve theorem


Counterexamples

2. For closed good covers some “finiteness” is needed:
 - ▶ Consider the cover of S^1 by its points
 - ▶ As the nerve $\text{Nrv } \mathcal{A}$ is a disjoint union of points, it is not homotopy equivalent to S^1
 - ▶ All conditions in 1.b) and 2.a) are satisfied except the locally finiteness assumption

Unified nerve theorem

Counterexamples

3. The “latching assumption” is not a proof artefact:
- ▶ Consider the *double comb space* C and denote the two combs by A_1 and A_2




- ▶ The nerve $\text{Nrv } \mathcal{A}$ is contractible, but C is not
- ▶ The pairs $(A_1, A_1 \cap A_2)$ and $(A_2, A_1 \cap A_2)$ do not satisfy the homotopy extension property

Unified nerve theorem

Counterexamples

3. The “latching assumption” is not a proof artefact; even if we only care about the homologies:
- ▶ Consider the *Warsaw circle* $W \subseteq S^2$ that separates the sphere into two connected components U_1 and U_2



- ▶ The closed sets $A_1 = U_1 \cup W$ and $A_2 = U_2 \cup W$ cover the sphere and are contractible
- ▶ $A_1 \cap A_2 = W$ is acyclic and hence $\mathcal{A} = \{A_1, A_2\}$ is a homologically good closed cover of S^2
- ▶ S^2 and $\text{Nrv } \mathcal{A}$ do not have isomorphic homology groups

Summary

- ▶ Gave proofs of (functorial) nerve theorems that are attractive to students and newcomers to TDA.
- ▶ Learned that functoriality depends on the framework and holds without any assumptions.
- ▶ Abstract homotopy theory can help to give a “unified view” on the functorial nerve theorem.

Future work:

- ▶ Discuss approximate nerve theorems.
- ▶ Discuss Latschev’s result on the reconstruction of Riemannian manifolds and its functoriality by using *Vietoris–Rips good covers*.

Funding: DFG (SFB/TRR 109 *Discretization in Geometry and Dynamics*)